

2008 BETTER HYBRIDS Harvest Report for Indiana South [INSO]

Keith Scott, Ripley County [VERSAILLES]

LATE SEASON TEST

113 - 116 Day CRM

SOIL CONDITIONS: Sandy clay loam, 0.5% slope, well drained, no-till
 1.0% o.m., 6.4 pH, moderate P & K

B8INSO17b

PREVIOUS CROP/HERB: Soybean / Roundup

FERTILITY PROGRAM: 180 lbs N as NH3 sidedress, 5 gal 6-24-6 as starter

PEST MANAGEMENT: Touchdown burndown fb Lexar
 Force 3G T-band

PLANTED - SEEDING: April 30 - 29,000 seed/A

TOP 30 OF 36 HYBRIDS FOR GROSS INCOME (SORTED BY YIELD)
HARVESTED - STAND:

AVERAGE of (3) REPLICATIONS

COMPANY	HYBRID	TECHNOLOGY†	IST††	YIELD Bu/A	MOIST %	LODGING %	STAND (x 1000)	GROSS INCOME
Location was lost due to heavy rain and cold temperatures after planting. Keith had to replant all of his corn.								
Test Average = Isd(.10) = C.V. =								

Yield & Income Factors: Base Moisture = 15.0% Shrink = 1.4 Drying = \$0.035 Price = \$2.20

 † RESISTANCE TECHNOLOGY: **Corn Borer** - CB, HX1, HXT, YGCB, YGPL, YGVT3; **Corn Rootworm** - HXT, RW, YGPL, YGRW, YGVT2, YGVT3; **Clearfield** - CL; **Glyphosate** - GT, RR2, YGVT2, YGVT3; **Liberty** - HX1, HXT, LL

 †† IST: **C250, C1250** - Cruiser @ 0.250 and 1.250 g ai/seed. **P250 and P1250** - Poncho @ 0.250 and 1.250 g ai/seed.

→ identifies the check hybrid entered in both early- and full-season tests to assist in comparing the results.

* indicates seed tested from lots not commercially available at planting. Hybrids in italics exceed the grain moisture limit for this test.

FIELD NOTES: